
# Mascia Brunelli *Hemostats Family*

CUTANPLAST®

CUTANPLAST® Fast

emosist®




**Easy to use**
***Facile da usare***

**Different sizes and format**
***Differenti formati***

**Easy to cut**
***Facile da tagliare***

**Ready to use, dry**
***Pronto all'uso***

**Better haemostasis management**
***Migliore controllo dell'emorragia***

**Fast fluid absorption**
***Veloce assorbimento dei liquidi***

**Oxidized regenerated cellulose**
***Tessuto in cellulosa ossidata rigenerata***

**Ready for minimally invasive procedure**
***Adatto ad utilizzo laparoscopio***

**Ready to suture and wrap up**
***Pronto da suturare e avvolgere***

Gelatin Sponges

CUTANPLAST®

CUTANPLAST® previously soaked with sterile saline is applied pressing lightly on the clean wound, until the adhesion exerted by fibrin is obtained. The sponge can be left in situ, or removed just after the hemostasis occurred. CUTANPLAST® is completely absorbed within 3 weeks approximately.

- Has a neutral pH;
- May be soaked with sterile physiological solution or with antibiotics;
- Many sizes and formats are avaliable
- May be easily reduced in size to any required format;
- Easy to use;
- Safe;
- Reliable
- Avaliable in POWDER FORMAT




CUTANPLAST Fast®

CUTANPLAST FAST® is an adjuvant in the control of local bleeding, in the event that normal surgical procedures are ineffective. The CUTANPLAST FAST® product absorbs in the first minute an amount of liquid equal to 60 times its own weight. Due to the high and rapid absorption capacity of liquids, CUTANPLAST FAST® is specifically indicated for all surgeries involving big organs.
Helps to manage different types of bleeding, thanks to the possibility of being soaked with pharmaceutical substances. The product is compact and flexible and it is suitable for laparoscopy and for biopsy procedures.

- High absorbency : up to 60 times its own weight in the first min
- Ready to use dry
- Haemostatic rates : 2 - 8 minutes
- Dissolution rates: 2 - 4 weeks
- Its helps reduce surgery time
- May be easily reduced in size to any required format
- Easy to use;
- Safe
- Avaliable in POWDER FORMAT



# Mascia Brunelli *Hemostats Family*

emosist®

Oxidized Regenerated Cellulose

EMOSIST® is made from oxidized regenerated cellulose (Viscose). The oxidation process of the rayon-viscose gives chemical and physical characteristics to the finished product that can denature the proteins of the blood quickly activating the process of blood clotting.


- Haemostatic rate: 2 - 8 minutes
- Dissolution rate: 1 - 2 weeks
- High biocompatibility
- the pH rate between 2 and 3 gives to the product bacteriostatic properties in vitro tested, the product must not be used as an alternative to therapeutic or prophylactic antimicrobial agents.
- High ductility of use both dry and soaked in blood, does not adhere to tools or to surgical gloves.
- Easy to drape, wrap or suture
- Ready for M.I.S.




## Product Properties


### Specialities Indications

H: High/Consistente
M: Medium/Medio
L: Low/Basso




### Products Key Features

1 = Very Low - 7 = Very High



## *Products Literatures:*

**Biocompatibility and performance in vitro of a hemostatic gelatin sponge.**
Journal of Biomaterials Science, Polymer Edition, 11:7, 685-699, 2000.
Elisabetta Cenni, Gabriela Ciapetti, Susanna Stea, Alessandra Corradini, Fiorenzo Carozzi.

**Comparative analysis of Cutanplast® and Spongostan nasal packing after endoscopic sinus surgery: a prospective, randomized, multicenter study.**
Eur Arch Otorhinolaryngol. 2015 Jul;272(7):1699-705. doi: 10.1007/s00405-014-3264-3.
Epub 2014 Sep 4.
Cho KS, Park CH, Hong SL, Kim MJ, Kim JY, Kim YW, Koo SK, Roh HJ.

**Effectiveness of hemostatic gelatin sponge as a packing material after septoplasty: a prospective, randomized, multicenter study.**
Auris Nasus Larynx. 2018 Apr;45(2):286-290. doi: 10.1016/j.anl.2017.05.007. Epub 2017 May 25.
Kim SD, Hong SL, Kim MJ, Kim JY, Kim YW, Koo SK, Cho KS.

**Use of oxidized regenerated cellulose to achieve hemostasis during laparoscopic cholecystectomy: a retrospective cohort analysis.**
BMC Res Notes (2018) 11:239
Emilia Masci, Giuseppe Faillace, Mauro Longoni.

**Comparison of the efficacy of five standard topical hemostats: a study in porcine liver and spleen models of surgical bleeding**
BMC Surgery (2020) 20:215
Valerio Antonio Paternò, Alessandro Bisin, Alessandro Addis.

**Ptotic Right Retro-renal Liver Lobe Injury During Supine Percutaneous Nephrolithotomy Managed by Hemostatic Sponge**
Indian Journal of Surgery (on line 2021)
Ibrahim Üntan, Volkan Sabur.

Indice di Rev. MMB01 F01-1-11/2021

CE 0483

**Mascia Brunelli S.p.A.**
V.le Monza, 272 - 20128 Milano (Italy)
tel +39 02 25209.1- fax +39 02 2576428
mktg@masciabrunelli.it
www.masciabrunelli.it
www.cutanplast.com
www.emosist.it